

A Study on Discrete Model of Three Species Syn-Eco-System with Limited Resources

(One Period Equilibrium States)

B. Hari Prasad

Department of Mathematics, Chaitanya Degee College, Hanamkonda, A.P., India
Email: sumathi_prasad73@yahoo.com

Abstract—In this paper, the system comprises of a commensal (S_1), two hosts S_2 and S_3 i.e., S_2 and S_3 both benefit S_1 , without getting themselves effected either positively or adversely. Further S_2 is a commensal of S_3 , S_3 is a host of both S_1 , S_2 and all the three species have limited resources. The basic equations for this model constitute as three first order non-linear ordinary difference equations. All possible equilibrium points are identified based on the model equations and criteria for their stability are discussed. Further the numerical solutions are computed for specific values of the various parameters and the initial conditions.

Index Terms—Commensal, Discrete model, Equilibrium state, Integer, Host, Species, Stable.

AMS Classification: 92D25, 92D40.

I. INTRODUCTION

Ecology, a branch evolutionary biology, deals with living species that coexist in a physical environment sustain themselves on common resources. It is a common observations that the species of same nature can not flourish is isolation without any interaction with species of different kinds. Syn-ecology is an ecosystem comprising of two or more distinct species. Species interact with each other in one way or other. The Ecological interactions can be broadly classified as Ammensalism, Competition, Commensalism, Neutralism, Mutualism, Predation Parasitism and so on. Lotka[1] and Volterra [2] pioneered theoretical ecology significantly and opened new eras in the field of life and biological sciences.

Mathematical Modeling is a vital role in providing insight in to the mutual relationships (positive, negative) between the interacting species. The general concepts of modeling have been discussed by several authors, viz., [3-6]. Srinivas[7] studied competitive ecosystem of two species and three species with limited and unlimited resources. Later, Laxminarayan et al [8] studied prey-predator ecological models with partial cover for the prey and alternate food for the predator. Stability analysis of competitive species was carried out by [9-10], while Reddy [11] investigated mutualism between two species.

Acharyulu et al [12-13] derived some productive results on various mathematical models of ecological Ammensalism with multifarious resources in the manifold directions. Further Kumar [14] studied some mathematical models of ecological commensalism. The present author Prasad et al [15-22] investigated on the stability of a three and four species syn-ecosystems. Recently, Din et al [23] discussed the basic concepts on stability analysis of discrete ecological models and Kiran et al [24] worked on discrete model of three species ecosystem by considering interactions like prey-predator and commensalism.

The present investigation is a discrete model of three species (S_1, S_2, S_3) syn-eco system with limited resources. The system comprises of a commensal (S_1), two hosts S_2 and S_3 . Further S_2 is a commensal of S_3 , S_3 is a host of both S_1 and S_2 . *Commensalism* is a symbiotic interaction between two populations where one population (S_1) gets benefit from (S_2) while the other (S_2) is neither harmed nor benefited due to the interaction with (S_1). The benefited species (S_1) is called the commensal and the other, the helping one (S_2) is called the host species. A common example is an animal using a tree for shelter-tree (host) does not get any benefit from the animal (commensal).

II. BASIC EQUATIONS OF THE MODEL

The model equations for the three species syn-ecosystem is given by the following system of first order non-linear difference equations employing the following notation.

A. Notation Adopted

$N_i(t)$: The population strength of S_i at time t , $i = 1, 2, 3$
 t : Time instant
 a_i : Natural growth rate of S_i , $i = 1, 2, 3$
 a_{ii} : Self inhibition coefficient of S_i , $i = 1, 2, 3$
 a_{12}, a_{13} : Interaction coefficients of S_1 due to S_2 and S_3
 a_{23} : Interaction coefficient of S_2 due to S_3

Further the variables N_1, N_2, N_3 are non-negative and the model parameters $a_1, a_2, a_3, a_{11}, a_{12}, a_{13}, a_{22}, a_{23}, a_{33}$ are assumed to be non-negative constants.

B. Basic equations

Consider the growth of the species during the time interval $(t, t + 1)$

- Equation for the first species (N_1):

$$N_1(t+1) = N_1(t) + a_1N_1(t) - a_{11}N_1^2(t) + a_{12}N_1(t)N_2(t) + a_{13}N_1(t)N_3(t) \quad (1)$$

- Equation for the second species (N_2):

$$N_2(t+1) = N_2(t) + a_2N_2(t) - a_{22}N_2^2(t) + a_{23}N_2(t)N_3(t) \quad (2)$$

- Equation for the third species (N_3):

$$N_3(t+1) = N_3(t) + a_3N_3(t) - a_{33}N_3^2(t) \quad (3)$$

- Species-growth equations in the discrete form:

Consider the nonlinear autonomous system of discrete equations

$$N_1(t+1) = \alpha_1N_1(t) - a_{11}N_1^2(t) + a_{12}N_1(t)N_2(t) + a_{13}N_1(t)N_3(t) \quad (4)$$

$$N_2(t+1) = \alpha_2N_2(t) - a_{22}N_2^2(t) + a_{23}N_2(t)N_3(t) \quad (5)$$

$$N_3(t+1) = \alpha_3N_3(t) - a_{33}N_3^2(t) \quad (6)$$

where $\alpha_i = a_i + 1, i = 1, 2, 3$

III. EQUILIBRIUM STATES

For a continuous model the equilibrium states are defined by $\frac{dN_i}{dt} = 0, i = 1, 2, 3$, the equilibrium states for a discrete model are defined in terms of the period of no growth. i.e, $N_i(t+r) = N_i(t), r = 1, 2, 3, \dots$, where r is the period of the equilibrium state.

Kapur J.N, Mathematical Modeling through Discrete Mathematics-Fascinating World of Mathematical Sciences, 1989, Volume-II, pp.71-80, Mathematical Science Trust Society, New Delhi.

- One period equilibrium states (Stage-I)

$$N_i(t+1) = N_i(t), i = 1, 2, 3$$

The system under investigation has fourteen equilibrium states given by

- (i) Fully washed out state

$$E_1 : \bar{N}_1 = 0, \bar{N}_2 = 0, \bar{N}_3 = 0$$

- (ii) States in which only one of the tree species is survives while the other two are not

$$E_2 : \bar{N}_1 = 0, \bar{N}_2 = 0, \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}, \text{ when } \alpha_3 > 1$$

$$E_3 : \bar{N}_1 = 0, \bar{N}_2 = \frac{\alpha_2 - 1}{a_{22}}, \bar{N}_3 = 0, \text{ when } \alpha_2 > 1$$

$$E_4 : \bar{N}_1 = \frac{\alpha_1 - 1}{a_{11}}, \bar{N}_2 = 0, \bar{N}_3 = 0, \text{ when } \alpha_1 > 1$$

- (iii) States in which only two of the tree species are survives while the other one is not

$$E_5 : \bar{N}_1 = 0, \bar{N}_2 = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_2 > 1$ and $\alpha_3 > 1$

$$E_6 : \bar{N}_1 = 0, \bar{N}_2 = a_{23} \left(\frac{\alpha_3 - 1}{a_{22}a_{33}} \right), \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_2 = 1$ and $\alpha_3 > 1$

$$E_7 : \bar{N}_1 = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], \bar{N}_2 = 0, \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_1 > 1$ and $\alpha_3 > 1$

$$E_8 : \bar{N}_1 = a_{13} \left(\frac{\alpha_3 - 1}{a_{11}a_{33}} \right), \bar{N}_2 = 0, \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_1 = 1$ and $\alpha_3 > 1$

$$E_9 : \bar{N}_1 = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} \left(\frac{\alpha_2 - 1}{a_{22}} \right) \right], \bar{N}_2 = \frac{\alpha_2 - 1}{a_{22}}, \bar{N}_3 = 0$$

when $\alpha_1 > 1$ and $\alpha_2 > 1$

$$E_{10} : \bar{N}_1 = a_{12} \left(\frac{\alpha_2 - 1}{a_{11}a_{22}} \right), \bar{N}_2 = \frac{\alpha_2 - 1}{a_{22}}, \bar{N}_3 = 0$$

when $\alpha_1 = 1$ and $\alpha_2 > 1$

(iv) The co-existent states (or) normal steady states

$$E_{11} : \bar{N}_1 = \frac{1}{a_{11}} \left\{ (\alpha_1 - 1) + \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\},$$

$$\bar{N}_2 = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}},$$

when α_1, α_2 and $\alpha_3 > 1$

$$E_{12} : \bar{N}_1 = \frac{1}{a_{11}} \left\{ \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\},$$

$$\bar{N}_2 = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_1 = 1$ and $\alpha_2, \alpha_3 > 1$

$$E_{13} : \bar{N}_1 = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$$

$$\bar{N}_2 = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right), \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_2 = 1$ and $\alpha_1, \alpha_3 > 1$

$$E_{14} : \bar{N}_1 = \frac{1}{a_{11}} \left[a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$$

$$\bar{N}_2 = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right), \bar{N}_3 = \frac{\alpha_3 - 1}{a_{33}}$$

when $\alpha_1, \alpha_2 = 1$ and $\alpha_3 > 1$

IV. STABILITY ANALYSIS

A. Stability of $E_1(0, 0, 0)$

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = 0$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = 0$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = 0$$

i.e, $N_i(t+r) = 0$, where r is an integer and $i = 1, 2, 3$

Hence, $E_1(0, 0, 0)$ is stable.

B. Stability of E_2

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = 0$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = 0$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_i(t+r) = 0, N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$ where r is an integer and $i = 1, 2$

Hence, E_2 is stable.

C. Stability of E_3

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = 0$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = \frac{\alpha_2 - 1}{a_{22}}$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = 0$$

i.e, $N_i(t+r) = 0, N_2(t+r) = \frac{\alpha_2 - 1}{a_{22}}$ where r is an integer

and $i = 1, 3$

Hence, E_3 is stable.

D. Stability of E_4

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = \frac{\alpha_1 - 1}{a_{11}}$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = 0$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = 0$$

i.e, $N_1(t+r) = \frac{\alpha_1 - 1}{a_{11}}, N_i(t+r) = 0$ where r is an integer

and $i = 2, 3$

Hence, E_4 is stable.

E. Stability of E_5

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = 0$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots$$

$$= \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_1(t+r) = 0, N_2(t+r) = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$

$N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$, where r is an integer

Hence, E_5 is stable.

F. Stability of E_6

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = 0$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = a_{23} \left(\frac{\alpha_3 - 1}{a_{22}a_{33}} \right)$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_1(t+r) = 0, N_2(t+r) = a_{23} \left(\frac{\alpha_3 - 1}{a_{22}a_{33}} \right),$

$$N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}, \text{ where } r \text{ is an integer}$$

Hence, E_6 is stable.

G. Stability of E_7

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = 0$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_1(t+r) = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$

$$N_2(t+r) = 0, N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}, \text{ where } r \text{ is an integer}$$

Hence, E_7 is stable.

H. Stability of E_8

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = a_{13} \left(\frac{\alpha_3 - 1}{a_{11}a_{33}} \right)$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = 0$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_1(t+r) = a_{13} \left(\frac{\alpha_3 - 1}{a_{11}a_{33}} \right), N_2(t+r) = 0,$

$$N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}, \text{ where } r \text{ is an integer}$$

Hence, E_8 is stable.

I. Stability of E_9

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} \left(\frac{\alpha_2 - 1}{a_{22}} \right) \right]$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = \frac{\alpha_2 - 1}{a_{22}}$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = 0$$

i.e, $N_1(t+r) = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} \left(\frac{\alpha_2 - 1}{a_{22}} \right) \right],$

$$N_2(t+r) = \frac{\alpha_2 - 1}{a_{22}}, N_3(t+r) = 0, \text{ where } r \text{ is an integer}$$

Hence, E_9 is stable.

J. Stability of E_{10}

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = a_{12} \left(\frac{\alpha_2 - 1}{a_{11}a_{22}} \right)$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = \frac{\alpha_2 - 1}{a_{22}}$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = 0$$

i.e, $N_1(t+r) = a_{12} \left(\frac{\alpha_2 - 1}{a_{11}a_{22}} \right), N_2(t+r) = \frac{\alpha_2 - 1}{a_{22}},$

$$N_3(t+r) = 0, \text{ where } r \text{ is an integer}$$

Hence, E_{10} is stable.

K. Stability of E_{11}

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = \frac{1}{a_{11}} \left\{ (\alpha_1 - 1) + \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\}$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

i.e, $N_1(t+r) = \frac{1}{a_{11}} \left\{ (\alpha_1 - 1) + \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\},$

$$N_2(t+r) = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$$

where r is an integer

Hence, E_{11} is stable.

L. Stability of E_{12}

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots = \frac{1}{a_{11}} \left\{ \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\}$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

$$\text{i.e., } N_1(t+r) = \frac{1}{a_{11}} \left\{ \frac{a_{12}}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right] + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right\},$$

$$N_2(t+r) = \frac{1}{a_{22}} \left[(\alpha_2 - 1) + a_{23} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right], N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$$

where r is an integer
Hence, E_{12} is stable.

M. Stability of E_{13}

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots$$

$$= \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right)$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

$$\text{i.e., } N_1(t+r) = \frac{1}{a_{11}} \left[(\alpha_1 - 1) + a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$$

$$N_2(t+r) = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right), N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$$

where r is an integer
Hence, E_{13} is stable.

N. Stability of E_{14}

$$N_1(t) = N_1(t+1) = N_1(t+2) = \dots$$

$$= \frac{1}{a_{11}} \left[a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right]$$

$$N_2(t) = N_2(t+1) = N_2(t+2) = \dots = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right)$$

$$N_3(t) = N_3(t+1) = N_3(t+2) = \dots = \frac{\alpha_3 - 1}{a_{33}}$$

$$N_1(t+r) = \frac{1}{a_{11}} \left[a_{12} a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right) + a_{13} \left(\frac{\alpha_3 - 1}{a_{33}} \right) \right],$$

$$N_2(t+r) = a_{23} \left(\frac{\alpha_3 - 1}{a_{22} a_{33}} \right), N_3(t+r) = \frac{\alpha_3 - 1}{a_{33}}$$

where r is an integer
Hence, E_{14} is stable.

At this stage all the fourteen equilibrium states are stable.

• Two period equilibrium states (Stage-II)

$$N_i(t+2) = N_i(t), \quad i = 1, 2, 3$$

At this stage the system has 125 equilibrium states. The present paper deals only the stability of one period equilibrium states and the stability of the two period equilibrium states will be presented in the forth coming communications.

V. NUMERICAL EXAMPLES

The numerical solutions of the discrete model equations computed for specific values of the various parameters and the initial conditions. The results are illustrated in Figures 1 to 6.




Fig.1. Variation of population against time for $\alpha_1=3.9$, $\alpha_2=2.6$, $\alpha_3=4.4$, $a_{11}=1.18$, $a_{12}=0.6$, $a_{13}=1.5$, $a_{22}=0.8$, $a_{23}=1.2$, $a_{33}=1.7$, $N_{10}=7.5$, $N_{20}=5$, $N_{30}=2$




Fig.2. Variation of population against time for $\alpha_2=1$, $\alpha_3=2.5$, $a_{22}=2.3$, $a_{23}=4.4$, $a_{33}=0.4$, $N_{10}=0$, $N_{20}=7.5$, $N_{30}=3.75$




Fig.3. Variation of population against time for $\alpha_2=1$, $\alpha_3=4.2$, $a_{11}=3.2$, $a_{13}=1.6$, $a_{33}=0.5$, $N_{10}=3.2$, $N_{20}=0$, $N_{30}=6.4$




Fig.4 Variation of population against time for $\alpha_2=1$, $\alpha_2=2.7$, $a_{11}=1.4$, $a_{12}=5.85$, $a_{22}=0.8$, $N_{10}=8.9$, $N_{20}=2.1$, $N_{30}=0$




Fig.5. Variation of population against time for $\alpha_1=3.8$, $a_{11}=0.6$, $N_{10}=4.7$, $N_{20}=0$, $N_{30}=0$




Fig.6. Variation of population against time for $N_{10}=0$, $N_{20}=0$, $N_{30}=0$

VI. CONCLUSION

The present paper deals with an investigation on a discrete model of three species syn eco-system with limited resources. The basic equations for this model constitute as three first order non-linear coupled ordinary difference equations. All possible equilibrium points of the model are identified at $N_i(t+1) = N_i(t)$; $i = 1, 2, 3$. It is observe that, all the fourteen equilibrium points are found to be stable. Further the numerical solutions for the discrete model equations are computed.

ACKNOWLEDGMENT

I thank to Professor (Retired), N.Ch. Pattabhi Ramacharyulu, Department of Mathematics, National Institute of Technology, Warangal (A.P.), India for his valuable suggestions and encouragement.

REFERENCES

- [1] A.J. Lotka, "Elements of Physical Biology", Williams and Wilking, Baltimore, 1925.
- [2] V. Volterra, "Leconsen La Theorie Mathematique De La Letite Pou Lavie", Gauthier-Villars, Paris, 1931.
- [3] A.P. Colinvaux, "Ecology", John Wiley, New York, 1986.
- [4] J.N. Kapur, "Mathematical Modeling in Biology and Medicine", Affiliated East West, 1985.
- [5] J.M. Kushing, "Integro-Differential Equations and Delay Models in Population Dynamics", Lecture Notes in Bio-Mathematics, Springer Verlag, 20, 1977.
- [6] W.J. Meyer, "Concepts of Mathematical Modeling", Mc.Grawhill, 1985.
- [7] N.C. Srinivas, "Some Mathematical Aspects of Modeling in Bio-medical Sciences", Kakatiya University, Ph.D Thesis, 1991.
- [8] K.L. Narayan and N.Ch.P.R. Charyulu, "A Prey-Predator Model with Cover for Prey and Alternate Food for the Predator and Time Delay", Int. J. of Scientific Computing, vol. 1, pp. 7 – 14, 2007.
- [9] R.A. Reddy, N.Ch.P.R. Charyulu and B.K. Gandhi, "A Stability Analysis of Two Competitive Interacting Species with Harvesting of Both the Species at a Constant Rate", Int. J. of Scientific Computing, vol. 1, pp. 57 – 68, 2007.
- [10] B. B. Rama Sharma and N.Ch. P.R. Charyulu, "Stability Analysis of Two Species Competitive Eco-system, Int. Journal of Logic Based Intelligent Systems, vol. 2, 2008.
- [11] R.R. Reddy, "A Study on Mathematical Models of Ecological Mutualism between Two Interacting Species", Osmania University, Ph.D. Thesis, 2008.
- [12] K.V.L.N. Acharyulu and N.Ch.P.R. Charyulu, "An Enemy- Ammensal Species Pair With Limited Resources –A Numerical Study", Int. J. Open Problems Compt. Math., vol. 3, pp. 339– 356, 2010.
- [13] K. V. L. N. Acharyulu and N. Ch. P. R. Charyulu, "An Ammensal-Prey with three species Ecosystem", Int. J. of Computational Cognition , vol. 9, pp. 30 – 39, 2011.
- [14] N.P. Kumar, "Some Mathematical Models of Ecological Commensalism", Acharya Nagarjuna University, Ph.D. Thesis, 2010.
- [15] B.H. Prasad and N.Ch.P.R. Charyulu, "On the Stability of a Four Species: A Prey-Predator-Host-Commensal-Syn Eco-System-VII", Int. J. of Applied Mathematical Analysis and Applications, vol. 6, pp. 85 – 94, 2011.
- [16] B.H. Prasad and N.Ch.P.R. Charyulu, "On the Stability of a Four Species Syn Eco-System with Commensal Prey Predator Pair with Prey Predator Pair of Hosts-VII", J. of Comm. and Computer, vol. 8, pp. 415 – 421, 2011.
- [17] B.H. Prasad and N.Ch.P.R. Charyulu, "On the Stability of a typical three species syn eco-system", Int. J. of Mathematical Archive, vol. 3, pp. 3583 – 3601, 2012.
- [18] B.H. Prasad and N.Ch.P.R. Charyulu, "On the Stability of a Four Species Syn Eco-System with Commensal Prey Predator Pair with Prey Predator Pair of Hosts-VI", Matematika, vol. 28, pp. 181 – 192, 1012.
- [19] B.H. Prasad and N.Ch.P.R. Charyulu, "Global Stability of Four Species Syn Eco-System: A Prey-Predator-Host-Commensal - A Numerical Approach", Int. Journal of

Mathematics and Computer Applications Research, vol. 3, pp. 275 – 290, 2013.

- [20] B.H. Prasad and N.Ch.P.R. Charyulu, “Discrete Model of Commensalism Between Two Species”, International Journal of Modern Education and Computer Science, vol. 8, pp. 40 – 46, 2012.
- [21] B.H. Prasad, “A Discrete Model of a Typical Three Species Syn- Eco – System with Unlimited Resources for the First and Third Species”, Asian Academic Research Journal of Multidisciplinary, vol. 1, pp. 36 – 46, 2014.
- [22] B.H. Prasad, “On the Stability of a Three Species Syn-Eco-System with Mortality Rate for the Second Species”, International Journal of Social Science & Interdisciplinary Research, vol. 3, pp. 35 – 45, 2014.
- [23] Q. Din and E.M. Elsayed, “Stability Analysis of a Discrete Ecological Model”, Computational Ecology and Software, vol. 4, pp. 89 – 103, 1014.
- [24] D.R. Kiran, B.R. Reddy and N.Ch.P.R. Charyulu, “Discrete Time Stability Analysis of a Three Species Eco-System”, Universal J. of Applied Mathematics, vol. 2, pp.161 – 169, 1014.

Authors' Profile



B. Hari Prasad: He works as an Assistant Professor, Department of Mathematics, Chaitanya Degree and PG College (Autonomous), Hanamkonda, Andhra Pradesh, India. He has obtained M.Phil in Mathematics. He has submitted Ph.D. thesis for the award of Doctorate degree by the Dravidian University, Kuppam. His research papers in area of Applied Mathematics are more than **30** were published in various esteemed National and International Journals. He has zeal to find out new vistas in Mathematics.

How to cite this paper: B. Hari Prasad, "A Study on Discrete Model of Three Species Syn-Eco-System with Limited Resources — (One Period Equilibrium States)", IJMECS, vol.6, no.11, pp.38-44, 2014.DOI: 10.5815/ijmeecs.2014.11.05